

## Calorimetry and Specific Heat Calculation Worksheet

1. Find the equilibrium temperature if 200.0 grams of cold water at 10.0°C is mixed with 600.0 grams of warm water at 60.0°C.
2. 350 g of ice at 0.00°C are added to 50.0 g of steam at 140°C. Find  $T_f$ .
3. A 600 ml sample of water is at 80.0°C. How many grams of ice at -20.0°C must be added to bring the temperature down to 5.00°C?
4. An iron skillet weighing 1300. g is heated on a stove to 178.° C. How many joules have to be removed in order for the skillet to cool to 21.° C? ( $c_{Fe} = 0.449 \text{ J/g-}^\circ\text{C}$ )

5. When ice melts into liquid water at  $0.0^{\circ}\text{C}$ , it absorbs  $0.334\text{ kJ/g}$ . Suppose the amount of heat needed to melt  $38.0\text{ g}$  of ice is absorbed from  $0.210\text{ kg}$  of water contained in a glass, at a temperature of  $21.0^{\circ}\text{C}$ . What is the final temperature of the water in the glass?
6. A hot,  $80.0\text{-g}$  iron spoon ( $T=95.0^{\circ}\text{C}$ ) is placed in a coffee cup containing  $100.0\text{ g}$  of water at  $20^{\circ}\text{C}$ . After the spoon and the water reach thermal equilibrium, what is the temperature of the water? Use the following information: ( $C_{\text{Fe}}=0.45\text{ J/g K}$ ,  $C_{\text{w}}=4.184\text{ J/g K}$ ).
7. A  $400.0\text{ g}$  sample of methanol at  $16.0^{\circ}\text{C}$  is mixed with  $400.0\text{ g}$  of water at  $85.0^{\circ}\text{C}$ . Assuming no heat loss to the surroundings, what is the final temperature of the mixture? The specific heat of methanol is  $2450\text{ J/kg}\cdot^{\circ}\text{C}$ .

8. A 1.0 kg sample of metal with a specific heat of 0.50 KJ/KgC is heated to 100.0C and then placed in a 50.0 g sample of water at 20.0C. What is the final temperature of the metal and the water?
9. A 2.50 g sample of zinc is heated, then placed in a calorimeter containing 65.0 g of water. Temperature of water increases from 20.00 °C to 22.50 °C. The specific heat of zinc is 0.390 J/g°C. What was the initial temperature of the zinc metal sample? (final temperatures of zinc and water are the same)
10. A 13.5 g sample of gold is heated, then placed in a calorimeter containing 60.0 g of water. Temperature of water increases from 19.00 °C to 20.00 °C. The specific heat of gold is 0.130 J/g°C. What was the initial temperature of the gold metal sample?

11. A 28.4 g sample of aluminum is heated to 39.4 °C, then is placed in a calorimeter containing 50.0 g of water. Temperature of water increases from 21.00 °C to 23.00 °C. What is the specific heat of aluminum?
12. A 25.0g sample of an alloy was heated to 100 °C and dropped into a beaker containing 90 grams of water at 25.32 °C. The temperature of the water rose to a final temperature of 27.18°C. Neglecting heat losses to the room and the heat capacity of the beaker itself, what is the specific heat of the alloy?
13. Phileas Fogg, the character who went around the world in 80 days, was very fussy about his bathwater temperature. It had to be exactly 38.0° C. You are his butler, and one morning while checking his bath temperature, you notice that it's 42.0°C. You plan to cool the 100.0 kg of water to the desired temperature by adding an aluminum-duckie originally at freezer temperature (-24.0°C). Of what mass should the Al-duckie be? [Specific heat of Al = 0.900 J/(g°C); density of water = 1.00 g/ml]. Assume that no heat is lost to the air.

14. A hot cup of tea is often a beverage of choice in the winter because it warms us up. The temperature of the tea is  $50.0^{\circ}\text{C}$  and the normal body temperature is  $37.2^{\circ}\text{C}$ . If you drink a cup of tea with a mass of 256.3 grams, how much heat energy was transferred to your body?  $C_p$  of water =  $4.184 \text{ J/g}^{\circ}\text{C}$
15. You want to do an experiment to measure the conversion of gravitational potential energy to kinetic energy to heat by dropping 2.0 kg of copper off the roof of LEHS, a height of 14 m. How much will the temperature of the copper increase?
16. Based on your answer to question #5 above, you decide to modify your experiment by dropping the 2.0kg bag of copper from a height of 2.0m to the floor multiple times. How many times would you need to drop the copper bag to get a temperature increase of  $2^{\circ}\text{C}$ ?

## Heat of Vaporization and Heat of Fusion Worksheet

1. How much heat is required to melt 360 g of solid water? Important constant:  $H_{\text{fus}}$  of water is 334 J/g.
2. How much heat is required to vaporized 24 g of liquid water?  $H_{\text{vap}}$  of water is 2257 J/g.
3. For a 500g block of lead (Pb) to melt, how much energy is needed?  $H_{\text{fus}}$  of lead is 23 J/g.
4. Mercury is a metal that is a liquid at room temperature. In order to solidify 7.5g of it, how much energy needs to be removed?  $H_{\text{fus}}$  of Hg is 11.3 J/g. 5. What is the heat of vaporization of ammonia ( $\text{NH}_3$ ) if 0.15 g of it requires 206.5 J for evaporation?
5. Ethanol ( $\text{C}_2\text{H}_5\text{OH}$ ) very easily changes from a liquid to a gas. If 29.34g of ethanol uses 32.23J of energy what would its  $H_{\text{vap}}$  be?

6. Iron is the heaviest metal vaporized in the sun. Its  $H_{\text{vap}}$  is 6071.43 J/g. How much heat is needed to turn (keep) 0.5 kilograms of iron into a gas?
7. In order for 5g liquid Hydrogen to become a solid, 12J of energy must be removed. What is the  $H_{\text{fus}}$  for the element, Hydrogen?
8. If 2083 Joules are used to melt 5.26 grams of aluminum, what is the heat of fusion of aluminum?
9. What is the mass of a sample of Nickel, which completely melts after 3120 J of heat? ( $\Delta H_{\text{fus}}$  of nickel is 298 J/g)